




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/54, 9/12, C12Q 1/48, G01N 33/573</b>		A1	(11) International Publication Number: <b>WO 96/37619</b>
			(43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: <b>PCT/IT96/00106</b>		(81) Designated States: AU, BR, CA, CN, JP, RU, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: <b>24 May 1996 (24.05.96)</b>			
(30) Priority Data: <b>RM95A000343 25 May 1995 (25.05.95) IT</b>		Published With international search report.	
(71) Applicant (for all designated States except US): ISTITUTO DI RICERCHE DI BIOLOGIA MOLECOLARE P. ANGELETTI S.P.A. [IT/IT]; Via Pontina Km. 30.600, I-00040 Pomezia (IT).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): DE FRANCESCO, Raffaele [IT/IT]; Via Devich, 26, I-00146 Roma (IT). TOMEI, Licia [IT/IT]; Via Gadda, 173, I-00143 Roma (IT). BEHRENS, Sven-Erik [DE/DE]; Steinweg 2, D-35096 Weimar (DE).			
(74) Agents: DI CERBO, Mario et al.; Società Italiana Brevetti S.p.A., Piazza di Pietra, 39, I-00186 Roma (IT).			

(54) Title: METHOD FOR REPRODUCING IN VITRO THE RNA-DEPENDENT RNA POLYMERASE AND TERMINAL NUCLEOTIDYL TRANSFERASE ACTIVITIES ENCODED BY HEPATITIS C VIRUS (HCV)

## (57) Abstract

This is a method for reproducing in vitro the RNA-dependent RNA polymerase activity associated with hepatitis C virus. The method is characterized in that sequences contained in NSSB are used in the reaction mixture. The terminal nucleotidyl transferase activity, a further property of the NSSB protein, can also be reproduced using this method. The method takes advantage of the fact that the NSSB protein, either purified to apparent homogeneity or present in extracts of overproducing organisms, can catalyse the addition of ribonucleotides to the 3'-termini of exogenous or endogenous RNA molecules. The invention also relates to a composition of matter that comprises sequences contained in NSSB, and to the use of these compositions for the set up of an enzymatic test capable of selecting, for therapeutic purposes, compounds that inhibit the enzymatic activity associated with NSSB. The figure shows plasmids used in the method to produce hepatitis C virus RNA-dependent RNA polymerase and terminal nucleotidyl transferase in cultivated eukaryotic and prokaryotic cells.



P ETL = promoter of the gene coding for the PCNA protein

P PH = promoter of the polyhedrin gene

Amp = gene coding for the β-lactamase enzyme (ampicillin resistance)

LacZ (β-gal) = gene coding for the β-galactosidase enzyme

Col E1 = pBR322 replication origin

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD FOR REPRODUCING IN VITRO THE RNA-DEPENDENT RNA  
POLYMERASE AND TERMINAL NUCLEOTIDYL TRANSFERASE  
ACTIVITIES ENCODED BY HEPATITIS C VIRUS (HCV)

DESCRIPTION

5           The present invention relates to the molecular  
biology and virology of the hepatitis C virus (HCV).  
More specifically, this invention has as its object the  
RNA-dependent RNA polymerase (RdRp) and the nucleotidyl  
terminal transferase (TNTase) activities produced by HCV,  
10       methods of expression of the HCV RdRp and TNTase, methods  
for assaying in vitro the RdRp and TNTase activities  
encoded by HCV in order to identify, for therapeutic  
purposes, compounds that inhibit these enzymatic  
activities and therefore might interfere with the  
15       replication of the HCV virus.

As is known, the hepatitis C virus (HCV) is the main  
etiological agent of non-A, non-B hepatitis (NANB). It  
is estimated that HCV causes at least 90% of post-  
transfusional NANB viral hepatitis and 50% of sporadic  
20       NANB hepatitis. Although great progress has been made in  
the selection of blood donors and in the immunological  
characterization of blood used for transfusions, there is  
still a high number of HCV infections among those  
receiving blood transfusions (one million or more  
25       infections every year throughout the world).  
Approximately 50% of HCV-infected individuals develop  
cirrhosis of the liver within a period that can range  
from 5 to 40 years. Furthermore, recent clinical studies  
suggest that there is a correlation between chronic HCV  
30       infection and the development of hepatocellular  
carcinoma.

HCV is an enveloped virus containing an RNA positive  
genome of approximately 9.4 kb. This virus is a member  
of the Flaviviridae family, the other members of which are  
35       the flaviviruses and the pestiviruses. The RNA genome of  
HCV has recently been mapped. Comparison of sequences  
from the HCV genomes isolated in various parts of the

world has shown that these sequences can be extremely heterogeneous. The majority of the HCV genome is occupied by an open reading frame (ORF) that can vary between 9030 and 9099 nucleotides. This ORF codes for a single viral polyprotein, the length of which can vary from 3010 to 3033 amino acids. During the viral infection cycle, the polyprotein is proteolytically processed into the individual gene products necessary for replication of the virus. The genes coding for HCV structural proteins are located at the 5'-end of the ORF, whereas the region coding for the non-structural proteins occupies the rest of the ORF.

The structural proteins consist of C (core, 21 kDa), E1 (envelope, gp37) and E2 (NS1, gp61). C is a non-glycosylated protein of 21 kDa which probably forms the viral nucleocapsid. The protein E1 is a glycoprotein of approximately 37 kDa, which is believed to be a structural protein for the outer viral envelope. E2, another membrane glycoprotein of 61 kDa, is probably a second structural protein in the outer envelope of the virus.

The non-structural region starts with NS2 (p24), a hydrophobic protein of 24 kDa whose function is unknown. NS3, a protein of 68 kDa which follows NS2 in the polyprotein, is predicted to have two functional domains: a serine protease domain in the first 200 amino-terminal amino acids, and an RNA-dependent ATPase domain at the carboxy terminus. The gene region corresponding to NS4 codes for NS4A (p6) and NS4B (p26), two hydrophobic proteins of 6 and 26 kDa, respectively, whose functions have not yet been clarified. The gene corresponding to NS5 also codes for two proteins, NS5A (p56) and NS5B (p65), of 56 and 65 kDa, respectively.

Various molecular biological studies indicate that the signal peptidase, a protease associated with the endoplasmic reticulum of the host cell, is responsible for proteolytic processing in the non-structural region,

that is to say at sites C/E1, E1/E2 and E2/NS2. A virally-encoded protease activity of HCV appears to be responsible for the cleavage between NS2 and NS3. This protease activity is contained in a region comprising  
5 both part of NS2 and the part of NS3 containing the serine protease domain, but does not use the same catalytic mechanism. The serine protease contained in NS3 is responsible for cleavage at the junctions between S3 and NS4A, between NS4A and NS4B, between NS4B and NS5A  
10 and between NS5A and NS5B.

Similarly to other (+)-strand RNA viruses, the replication of HCV is thought to proceed via the initial synthesis of a complementary (-)-RNA strand, which serves, in turn, as template for the production of  
15 progeny (+)-strand RNA molecules. An RNA-dependent RNA polymerase (RdRp) has been postulated to be involved in both these steps. An amino acid sequence present in all the RNA-dependent RNA polymerases can be recognized within the NS5 region. This suggests that the NS5 region  
20 contains components of the viral replication machinery. Virally-encoded polymerases have traditionally been considered important targets for inhibition by antiviral compounds. In the specific case of HCV, the search for such substances has, however, been severely hindered by  
25 the lack of both a suitable model system of viral infection (e.g. infection of cells in culture or a facile animal model), and a functional RdRp enzymatic assay.

It has now been unexpectedly found that this important limitation can be overcome by adopting the  
30 method according to the present invention, which also gives additional advantages that will be evident from the following.

The present invention has as its object a method for reproducing in vitro the RNA-dependent RNA polymerase  
35 activity of HCV that makes use of sequences contained in the HCV NS5B protein. The terminal nucleotidyl transferase activity, a further property of the NS5B

protein, can also be reproduced using this method. The method takes advantage of the fact that the proteins containing sequences of NS5B can be expressed in either eukaryotic or prokaryotic heterologous systems: the recombinant proteins containing sequences of NS5B, either purified to apparent homogeneity or present in extracts of overproducing organisms, can catalyse the addition of ribonucleotides to the 3'-termini of exogenous RNA molecules, either in a template-dependent (RdRp) or template-independent (TNTase) fashion.

The invention also extends to a new composition of matter, characterized in that it comprises proteins whose sequences are described in SEQ ID NO: 1 or sequences contained therein or derived therefrom. It is understood that this sequence may vary in different HCV isolates, as all the RNA viruses show a high degree of variability. This new composition of matter has the RdRp activity necessary to the HCV virus in order to replicate its genome.

The present invention also has as its object the use of this composition of matter in order to prepare an enzymatic assay capable of identifying, for therapeutic purposes, compounds that inhibit the enzymatic activities associated with NS5B, including inhibitors of the RdRp and that of the TNTase.

Up to this point a general description has been given of the present invention. With the aid of the following examples, a more detailed description of specific embodiments thereof will now be given, in order to give a clearer understanding of its objects, characteristics, advantages and method of operation.

Figure 1 shows the plasmids constructs used for the transfer of HCV cDNA into a baculovirus expression vector.

Figure 2 shows the plasmids used for the in vitro synthesis of the D-RNA substrate of the HCV RNA-dependent RNA polymerase [pT7-7(DCoH)], and for the expression of

the HCV RNA-dependent RNA polymerase in *E. coli* cells [pT7-7(NS5B)], respectively.

Figure 3 shows a schematic drawing of (+) and (-) strands of D-RNA. The transcript contains the coding region of the DCoH mRNA. The DNA-oligonucleotides a, b and c were designed to anneal with the newly-synthesized antisense RNA and the DNA/RNA hybrid was subjected to cleavage with RNase H. The lower part of the scheme depicts the expected RNA fragment sizes generated by RNase digestion of the RNA (-) hybrid with oligonucleotides a, b and c, respectively.

#### DEPOSITS

*E. Coli* DH1 bacteria, transformed using the plasmids pBac 5B, pBac 25, pT7.7 DCoH and pT7.7NS5B - containing SEQ ID NO:1; SEQ ID NO:2; the cDNA for transcription of SEQ ID NO:12; and SEQ ID NO:1, respectively, filed on May 9, 1995 with The National Collections of Industrial and Marine Bacteria Ltd. (NCIMB), Aberdeen, Scotland, UK. under access numbers NCIMB 40727, 40728, 40729 and 40730, respectively.

#### EXAMPLE 1

Method of expression of HCV RdRp/TNTase in *Spodoptera frugiperda* clone 9 (Sf9) cultured cells.

Systems for expression of foreign genes in insect cultured cells, such as *Spodoptera frugiperda* clone 9 (Sf9) cells infected with baculovirus vectors are known in the art (V. A. Luckow, Baculovirus systems for the expression of human gene products, (1993) Current Opinion in Biotechnology 4, pp. 564-572). Heterologous genes are usually placed under the control of the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus of the *Bombix mori* nuclear polyhedrosis virus. Methods for the introduction of heterologous DNA in the desired site in the baculoviral vectors by homologous recombination are also known in the art (D.R. O'Reilly, L. K. Miller, V.A. Luckow, (1992),

Baculovirus Expression Vectors-A Laboratory Manual, W. H. Freeman and Company, New York).

5 Plasmid vectors pBac5B and pBac25 are derivatives of a derivative of pBlueBacIII (Invitrogen) and were constructed for transfer of genes coding for NS4B and other non-structural HCV proteins in baculovirus expression vectors. The plasmids are schematically illustrated in figure 1 and their construction is described in detail in Example 8. Selected fragments of  
10 the cDNA corresponding to the genome of the HCV-BK isolate (HCV-BK; Takamizawa, A., Mori, C., Fuke, I., Manabe, S., Murakami, S., Fujita, J., Onishi, E., Andoh, T., Yoshida, I. and Okayama, H., (1991) Structure and Organization of the Hepatitis C Virus Genome Isolated from Human Carriers *J. Virol.*, 65, 1105-1113) were cloned  
15 under the strong polyhedrin promoter of the nuclear polyhedrosis virus and flanked by sequences that allowed homologous recombination in a baculovirus vector.

In order to construct pBac5B, a PCR product  
20 containing the cDNA region encoding amino acids 2420 to 3010 of the HCV polyprotein and corresponding to the NS5B protein (SEQ ID NO:1) was cloned between the *Bam*HI and *Hind*III sites of pBlue BacIII. The PCR sense oligonucleotide contained a translation initiation  
25 signal, whereas the original HCV termination codon serves for translation termination.

pBac25 is a derivative of pBlueBacIII (Invitrogen) where the cDNA region coding for amino acids 810 to 3010 of the HCV-BK polyprotein (SEQ ID NO:2) was cloned  
30 between the *Nco*I and the *Hind*III restriction sites.

*Spodoptera frugiperda* clone 9 (Sf9) cells and baculovirus recombination kits were purchased from  
Invitrogen. Cells were grown on dishes or in suspension at 27°C in complete Grace's insect medium (Gibco)  
35 containing 10% foetal bovine serum (Gibco). Transfection, recombination, and selection of baculovirus constructs were performed as recommended by the



manufacturer. Two recombinant baculovirus clones, Bac25 and Bac5B, were isolated that contained the desired HCV cDNA.

For protein expression, Sf9 cells were infected either with the recombinant baculovirus Bac25 or Bac5B at a density of  $2 \times 10^6$  cells per ml in a ratio of about 5 virus particles per cell. 48-72 hours after infection, the Sf9 cells were pelleted, washed once with phosphate buffered saline (PBS) and carefully resuspended ( $7.5 \times 10^7$  cells per ml) in buffer A (10 mM Tris/Cl pH 8, 1.5 mM  $MgCl_2$ , 10 mM NaCl) containing 1 mM dithiothreitol (DTT), 1 mM phenylmethanesulphonyl-fluoride (PMSF, Sigma) and 4 mg/ml leupeptin. All the following steps were performed on ice: after swelling for 30 minutes, the cells were disrupted by 20 strokes in a Dounce homogeniser using a tight-fitting pestle. Glycerol, as well as the detergents Nonidet P-40 (NP40) and 3-[(3-Cholamidopropyl)-dimethyl-ammonio]-1-propanesulfonate (CHAPS), were added to final concentrations of 10% (v/v), 1% (v/v) and 0.5% (w/v), respectively, and the cellular extract was incubated for a further hour on ice with occasional agitation. The nuclei were pelleted by centrifugation for 10 minutes at  $1000 \times g$ , and the supernatant was collected. The pellet was resuspended in buffer A containing the above concentrations of glycerol and detergents (0.5 ml per  $7.5 \times 10^7$  nuclei) by 20 strokes in the Dounce homogeniser and then incubated for one hour on ice. After repelleting the nuclei, both supernatants were combined, centrifuged for 10 minutes at  $8000 \times g$  and the pellet was discarded. The resulting crude cytoplasmic extract was used either directly to determine the RdRp activity or further purified on a sucrose gradient (see Example 5).

Infection of Sf9 cells with either the recombinant baculovirus Bac25 or Bac5B leads to the expression of the expected HCV proteins. Indeed, following infection of Sf9 cells with Bac25, correctly-processed HCV NS2 (24

kDa), NS3 (68 kDa), NS4B (26 kDa), NS4A (6 kDa), NS5A (56 kDa) and NS5B (65 kDa) proteins can be detected in the cell lysates by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and immunostaining. Following infection of  
5 Sf9 cells with Bac5B, only one HCV-encoded protein, corresponding in size to authentic NS5B (65 kDa), is detected by SDS-PAGE followed by immuno- or Coomassie Blue staining.

#### EXAMPLE 2

#### 10 Method of assay of recombinant HCV RdRp on a synthetic RNA template/substrate.

The RdRp assay is based on the detection of labelled nucleotides incorporated into novel RNA products. The *in vitro* assay to determine RdRp activity was performed in a  
15 total volume of 40 µl containing 1-5 µl of either Sf9 crude cytoplasmic extract or purified protein fraction. Unfractionated or purified cytoplasmic extracts of Sf9 cells infected with Bac25 or Bac5B may be used as the source of HCV RdRp. A Sf9 cell extract obtained from  
20 cells infected with a recombinant baculovirus construct expressing a protein that is not related to HCV may be used as a negative control. The following supplements are added to the reaction mixture (final concentrations): 20 mM Tris/Cl pH 7.5, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 25 mM KCl, 1  
25 mM EDTA, 5-10 µCi [<sup>32</sup>P] NTP of one species (unless otherwise specified, GTP, 3000 Ci/mmol, Amersham, was used), 0.5 mM each NTP (i.e. CTP, UTP, ATP unless specified otherwise), 20 U RNasin (Promega), 0.5 µg RNA-substrate (ca. 4 pmol; final concentration 100 nM), 2 µg  
30 actinomycin D (Sigma). The reaction was incubated for two hours at room temperature, stopped by the addition of an equal volume of 2 x Proteinase K (PK, Boehringer Mannheim) buffer (300 mM NaCl, 100 mM Tris/Cl pH 7.5, 1% w/v SDS) and followed by half an hour of treatment with  
35 50 µg of PK at 37°C. RNA products were PCA extracted, precipitated with ethanol and analysed by electrophoresis on 5% polyacrylamide gels containing 7M urea.

The RNA substrate we normally used for the assay (D-RNA) had the sequence reported in SEQ ID NO: 12, and was typically obtained by *in vitro* transcription of the linearized plasmid pT7-7(DCoH) with T7 polymerase, as described below.

Plasmid pT7-7(DCoH) (figure 2) was linearized with the unique *Bgl*II restriction site contained at the end of the DCoH coding sequence and transcribed *in vitro* with T7 polymerase (Stratagene) using the procedure described by the manufacturer. Transcription was stopped by the addition of 5 U/10 $\mu$ l of DNaseI (Promega). The mixture was incubated for a further 15 minutes and extracted with phenol/chloroform/ isoamylalcohol (PCA). Unincorporated nucleotides were removed by gel-filtration through a 1-ml Sephadex G50 spun column. After extraction with PCA and ethanol precipitation, the RNA was dried, redissolved in water and its concentration determined by optical density at 260 nm.

As will be clear from the experiments described below, any other RNA molecule other than D-RNA, may be used for the RdRp assay of the invention.

The above described HCV RdRp assay gave rise to a characteristic pattern of radioactively-labelled reaction products: one labelled product, which comigrated with the substrate RNA was observed in all reactions, including the negative control. This RNA species could also be visualised by silver staining and was thus thought to correspond to the input substrate RNA, labelled most likely by terminal nucleotidyl transferase activities present in cytoplasmic extracts of baculovirus-infected Sf9 cells. In the reactions carried out with the cytoplasmic extracts of Sf9 cells infected with either Bac25 or Bac5B, but not of cells infected with a recombinant baculovirus construct expressing a protein that is not related to HCV, an additional band was observed, migrating faster than the substrate RNA. This latter reaction product was found to be labelled to a

high specific activity, since it could be detected solely by autoradiography and not by silver staining. This novel product was found to be derived from the externally-added RNA template, as it was absent from control reactions where no RNA was added. Interestingly, the formation of a labelled species migrating faster than the substrate RNA was consistently observed with a variety of template RNA molecules, whether containing the HCV 3'-untranslated region or not. The 399 nucleotide mRNA of the liver-specific transcription cofactor DCoH (D-RNA) turned out to be an efficiently accepted substrate in our RdRp assay.

In order to define the nature of the novel species generated in the reaction by the Bac25- or Bac5B-infected cell extracts, we carried out the following series of experiments. (i) The product mixture was treated with RNase A or Nuclease P1. As this resulted in the complete disappearance of the radioactive bands, we concluded that both the labelled products were RNA molecules. (ii) Omission from the reaction mixtures of any of the four nucleotide triphosphates resulted in labelling of only the input RNA, suggesting that the faster migrating species is a product of a polymerisation reaction. (iii) Omission of  $Mg^{2+}$  ions from the assay caused a complete block of the reaction: neither synthesis of the novel RNA nor labelling of the input RNA were observed. (iv) When the assay was carried out with a radioactively labelled input RNA and unlabelled nucleotides, the labelled product was indistinguishable from that obtained under the standard conditions. We concluded from this result that the novel RNA product is generated from the original input RNA molecule.

Taken together, our data demonstrate that the extracts of Bac25- or Bac5B-infected Sf9 cells contain a novel magnesium-dependent enzymatic activity that catalyses *de novo* RNA synthesis. This activity was shown to be dependent on the presence of added RNA, but

independent of an added primer or of the origin of the input RNA molecule. Moreover, as the products generated by extracts of Sf9 cells infected with either Bac25 or Bac5B appeared to be identical, the experiments just described indicate that the observed RdRp activity is encoded by the HCV NS5B protein.

### EXAMPLE 3

#### Methods for the characterization of the HCV RdRp RNA product

The following methods were employed in order to elucidate the structural features of the newly-synthesized RNA product. Under our standard electrophoresis conditions (5% polyacrylamide, 7M urea), the size of the novel RNA product appeared to be approximately 200 nucleotides. This could be due to either internal initiation of RNA transcription, or to premature termination. These possibilities, however, appeared to be very unlikely, since products derived from RdRp assays using different RNA substrates were all found to migrate significantly faster than their respective templates. Increasing the temperature during electrophoresis and the concentration of acrylamide in the analytical gel lead to a significantly different migration behaviour of the RdRp product. Thus, using for instance a gel system containing 10% acrylamide, 7M urea, where separation was carried out at higher temperature, the RdRp product migrated slower than the input substrate RNA, at a position corresponding to at least double the length of the input RNA. A similar effect was observed when RNA-denaturing agents such as methylhydroxy-mercury ( $\text{CH}_3\text{HgOH}$ , 10 mM) were added to the RdRp products prior to electrophoresis on a low-percentage/lower temperature gel. These observations suggest that the RdRp product possesses an extensive secondary structure.

We investigated the susceptibility of the product molecule to a variety of ribonucleases of different specificity. The product was completely degraded upon

treatment with RNase A. On the other hand, it was found to be surprisingly resistant to single-strand specific nuclease RNase T1. The input RNA was completely degraded after 10 minutes incubation with 60 U RNase T1 at 22°C and silver staining of the same gel confirmed that not only the template, but also all other RNA usually detectable in the cytoplasmic extracts of Sf9 cells was completely hydrolysed during incubation with RNase T1. In contrast, the RdRp product remained unaltered and was affected only following prolonged incubation with RNase T1. Thus, after two hours of treatment with RNase T1, the labelled product molecule could no longer be detected at its original position in the gel. Instead, a new band appeared that had an electrophoretic mobility similar to the input template RNA. A similar effect was observed when carrying out the RNase T1 digestion for 1 hour, but at different temperatures: at 22°C, the RdRp product remained largely unaffected whereas at 37°C it was converted to the new product that co-migrates with the original substrate.

The explanation for these observations is that the input RNA serves as a template for the HCV RdRp, where the 3'-OH is used to prime the synthesis of the complementary strand by a turn-or "copy-back" mechanism to give rise to a duplex RNA "hairpin" molecule, consisting of the sense (template) strand to which an antisense strand is covalently attached. Such a structure would explain the unusual electrophoretic mobility of the RdRp product on polyacrylamide gels as well as its high resistance to single-strand specific nucleases. The turn-around loop should not be base-paired and therefore ought to be accessible to the nucleases. Treatment with RNase T1 thus leads to the hydrolysis of the covalent link between the sense and antisense strands to yield a double-stranded RNA molecule. During denaturing gel electrophoresis the two strands become separated and only the newly-synthesized

antisense strand, which should be similar in length to the original RNA template, would remain detectable. This mechanism would appear rather likely, especially in view of the fact that this kind of product is generated by several other RNA polymerases *in vitro*.

The following experiment was designed in order to demonstrate that the RNA product labelled during the polymerase reaction and apparently released by RNase T1 treatment exhibits antisense orientation with respect to the input template. For this purpose, we synthesized oligodeoxyribonucleotides corresponding to three separate sequences of the input template RNA molecule (figure 2), oligonucleotide a, corresponding to nucleotides 170-195 of D-RNA (SEQ ID NO: 3); oligonucleotide b, complementary to nucleotides 286-309 (SEQ ID NO: 4); oligonucleotide c, complementary to nucleotides 331-354 (SEQ ID NO: 5). These were used to generate DNA/RNA hybrids with the product of the polymerase reaction, such that they could be subjected to RNase H digests. Initially, the complete RdRp product was used in the hybridizations. However, as this structure is too thermostable, no specific hybrids were formed. The hairpin RNA was therefore pre-treated with RNase T1, denatured by boiling for 5 minutes and then allowed to cool down to room temperature in the presence of the respective oligonucleotide. As expected, exposure of the hybrids to RNase H yielded specific cleavage products. Oligonucleotide a-directed cleavage lead to products of about 170 and 220 nucleotides in length, oligonucleotide b yielded products of about 290 and 110 nucleotides and oligonucleotide c gave rise to fragments of about 330 and 65 nucleotides. As these fragments have the expected sizes (see figure 3), the results indicate that the HCV NS5B-mediated RNA synthesis proceeds by a copy-back mechanism that generates a hairpin-like RNA duplex.

#### EXAMPLE 4

Method of assay of recombinant HCV TNTase on a synthetic RNA substrate

The TNTase assay is based on the detection of template-independent incorporation of labelled nucleotides to the 3' hydroxyl group of RNA substrates. The RNA substrate for the assay (D-RNA) was typically obtained by *in vitro* transcription of the linearized plasmid pT7-7DCOH with T7 polymerase as described in Example 2. However, any other RNA molecule, other than D-RNA, may be used for the TNTase assay of the invention.

The *in vitro* assay to determine TNTase activity was performed in a total volume of 40 µl containing 1-5 µl of either Sf9 crude cytoplasmic extract or purified protein fraction. Unfractionated or purified cytoplasmic extracts of Sf9 cells infected with Bac25 or Bac5B may be used as the source of HCV TNTase. An Sf9 cell extract obtained from cells infected with a recombinant baculovirus construct expressing a protein that is not related to HCV may be used as a negative control. The following supplements are added to the reaction mixture (final concentrations): 20 mM Tris/Cl pH 7.5, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 25 mM KCl, 1 mM EDTA, 5-10 µCi [<sup>32</sup>P] NTP of one species (unless otherwise specified, UTP, 3000 Ci/mmol, Amersham, was used), 20 U RNasin (Promega), 0.5 µg RNA-substrate (ca. 4 pmol; final concentration 100 nM), 2 µg actinomycin D (Sigma). The reaction was incubated for two hours at room temperature, stopped by the addition of an equal volume of 2 x Proteinase K (PK, Boehringer Mannheim) buffer (300 mM NaCl, 100 mM Tris/Cl pH 7.5, 1% w/v SDS) and followed by half an hour of treatment with 50 µg of PK at 37°C. RNA products were PCA extracted, precipitated with ethanol and analysed by electrophoresis on 5% polyacrylamide gels containing 7M urea.

EXAMPLE 5

Method for the purification of the HCV RdRp/TNTase by sucrose gradient sedimentation



A linear 0.3-1.5 M sucrose gradient was prepared in buffer A containing detergents (see Example 1). Up to 2 ml of extract of Sf9 cells infected with Bac5B or Bac25 (corresponding to about  $8 \times 10^7$  cells) were loaded onto a 12 ml gradient. Centrifugation was carried out for 20 hours at 39000 x g using a Beckman SW40 rotor. 0.5 ml fractions were collected and assayed for activity. The NS5B protein, identified by western blotting, was found to migrate in the density gradients with an unexpectedly high sedimentation coefficient. The viral protein and ribosomes were found to co-sediment in the same gradient fractions. This unique behaviour enabled us to separate the viral protein from the main bulk of cytoplasmic proteins, which remained on the top of the gradient. The RdRp activity assay revealed that the RdRp activity co-sedimented with the NS5B protein. A terminal nucleotidyl transferase activity (TNTase) was also present in these fractions.

#### EXAMPLE 6

##### Method for the purification of the HCV TNTase/RdRp from Sf9 cells

Whole cell extracts are made from 1 g of Sf9 cells infected with Bac5B recombinant baculovirus. The frozen cells are thawed on ice in 10 ml of buffer containing 20 mM Tris/HCl pH 7.5, 1 mM EDTA, 10 mM DTT, 50% glycerol (N buffer) supplemented with 1 mM PMSF. Triton X-100 and NaCl are then added to a final concentration of 2% and 500 mM, respectively, in order to promote cell breakage. After the addition of  $MgCl_2$  (10 mM) and DNase I (15  $\mu g/ml$ ), the mixture is stirred at room temperature for 30 minutes. The extract is then cleared by ultracentrifugation in a Beckman centrifuge, using a 90 Ti rotor at 40,000 rpm for 30 minutes at 4° C. The cleared extract is diluted with a buffer containing 20 mM Tris/HCl pH 7.5, 1 mM EDTA, 10 mM DTT, 20% glycerol, 0.5% Triton X-100 (LG buffer) in order to adjust the NaCl concentration to 300 mM and incubated batchwise with 5 ml

of DEAE-Sepharose Fast Flow, equilibrated in LG buffer containing 300 mM NaCl. The matrix is then poured into a column and washed with two volumes of the same buffer. The flow-through and the first wash of the DEAE-Sepharose Fast Flow column is diluted 1:3 with LG buffer and applied onto a Heparin-Sepharose CL6B column (10 ml) equilibrated with LG buffer containing 100 mM NaCl. The Heparin-Sepharose CL6B is washed thoroughly and the bound proteins are eluted with a linear 100 ml gradient, from 100 mM to 1M NaCl in buffer LG. The fractions containing NS5B, as judged by silver- and immuno-staining of SDS-PAGE, are pooled and diluted with LG buffer in order to adjust the NaCl concentration to 50 mM. The diluted fractions are subsequently applied to a Mono Q-FPLC column (1 ml) equilibrated with LG buffer containing 50 mM NaCl. Proteins are eluted with a linear gradient (20 ml) from 50 mM to 1M NaCl in LG buffer. The fractions containing NS5B, as judged by silver- and immuno-staining of SDS-PAGE, are pooled and dialysed against LG buffer containing 100 mM NaCl. After extensive dialysis, the pooled fractions were loaded onto a PoyU-Sepharose CL6B (10 ml) equilibrated with LG buffer containing 100 mM NaCl. The PoyU-Sepharose CL6B was washed thoroughly and the bound proteins were eluted with a linear 100 ml gradient, from 100 mM to 1M NaCl in buffer LG. The fractions containing NS5B, as judged by silver- and immuno-staining of SDS-PAGE, are pooled, dialysed against LG buffer containing 100 mM NaCl and stored in liquid nitrogen prior to activity assay.

Fractions containing the purified protein NS5B were tested for the presence of both activities. The RdRp and TNTase activities were found in the same fractions. These results indicate that both activities, RNA-dependent RNA polymerase and terminal ribonucleotide transferase are the functions of the HCV NS5B protein.

We tested the purified NS5B for terminal nucleotidyl transferase activity with each of the four ribonucleotide

triphosphates at non-saturating substrate concentrations. The results clearly showed that UTP is the preferred TNTase substrate, followed by ATP, CTP and GTP irrespective of the origin of the input RNA.

5 EXAMPLE 7

Method of assay of recombinant HCV RdRp on a homopolymeric RNA template

Thus far we have described that HCV NS5B possesses an RNA-dependent RNA polymerase activity and that the  
10 synthesis of complementary RNA strand is a template-primed reaction. Interestingly, using unfractionated cytoplasmic extracts of Bac5B or Bac25 infected Sf9 cells as a source of RdRp we were not able to observe complementary strand RNA synthesis that utilized an  
15 exogenously added oligonucleotide as a primer. We reasoned that this could be due to the abundant ATP-dependent RNA-helicases that would certainly be present in our unfractionated extracts. We therefore wanted to address this question using the purified NS5B.

20 First of all, we wanted to establish whether the purified NS5B polymerase is capable of synthesizing RNA in a primer-dependent fashion on a homopolymeric RNA template: such a template should not be able to form intramolecular hairpins and therefore we expected that  
25 complementary strand RNA synthesis be strictly primer-dependent. We thus measured UMP incorporation dependent on poly(A) template and evaluated both oligo(rU)<sub>12</sub> and oligo(dT)<sub>12-18</sub> as primers for the polymerase reaction.

Incorporation of radioactive UMP was measured as follows.  
30 The standard reaction (10 -100 µl) was carried out in a buffer containing 20 mM Tris/HCl pH 7.5, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 25 mM KCl, 1 mM EDTA, 20 U RNasin (Promega), 1 µCi [<sup>32</sup>P] UTP (400 Ci/mmol, Amersham) or 1 µCi [<sup>3</sup>H] UTP (55 Ci/mmol, Amersham), 10 µM UTP, and 10 µg/ml poly(A) or  
35 poly(A)/oligo(dT)<sub>12-18</sub>. Oligo(U)<sub>12</sub> (1µg/ml) was added a primer. Poly A and polyA/oligodT<sub>12-18</sub> were purchased from Pharmacia. Oligo(U)<sub>12</sub> was obtained from Genset. The final

NS5B enzyme concentration was 10-100 nM. Under these conditions the reaction proceeded linearly for up to 3 hours. After 2 hours of incubation at 22°, the reaction was stopped by applying the samples to DE81 filters (Whatman), the filters washed thoroughly with 1M Na<sub>2</sub>HPO<sub>4</sub>/NaH<sub>2</sub>PO<sub>4</sub>, pH 7.0, rinsed with water, air dried and finally the filter-bound radioactivity was measured in a scintillation β-counter. Alternatively, the in vitro-synthesized radioactive product was precipitated by 10% trichloroacetic acid with 100 μg of carrier tRNA in 0.2 M sodium pyrophosphate, collected on 0.45-μm Whatman GF/C filters, vacuum dried, and counted in scintillation fluid.

Although some [<sup>32</sup>P]UMP or [<sup>3</sup>H]UMP incorporation was detectable even in the absence of a primer and is likely to be due to the terminal nucleotidyl transferase activity associated with our purified NS5B, up to 20% of product incorporation was observed only when oligo(rU)<sub>12</sub> was included as primer in the reaction mixture. Unexpectedly, also oligo(dT)<sub>12-18</sub> could function as a primer of poly(A)-dependent poly(U) synthesis, albeit with a lower efficiency.

Other template/primers suitable for measuring the RdRp activity of NS5B include poly(C)/oligo(G) or poly(C)/oligo(dG) in the presence of radioactive GTP, poly(G)/oligo(C) or poly(G)/oligo(dC) in the presence of radioactive CTP, poly(U)/oligo(A) or poly(U)/oligo(dA) in the presence of radioactive ATP, poly(I)/oligo(C) or poly(I)/oligo(dC) in the presence of radioactive CTP.

#### 30 EXAMPLE 8

##### Method of Expression Of HCV RdRp/TNTase in E. Coli

The plasmid pT7-7(NS5B), described in Figure 2 and Example 8, was constructed in order to allow expression in E. coli of the HCV protein fragment having the sequence reported in SEQ ID NO 1. Such protein fragment contains the RdRp and the TNTase of NS5B, as discussed above. The fragment of HCV cDNA coding for the NS5B

protein was thus cloned downstream of the bacteriophage T7  $\phi$ 10 promoter and in frame with the first ATG codon of the phage T7 gene 10 protein, using methods that are known to the molecular biology practice and described in detail in Example 8. The pT7-7(NS5B) plasmid also contains the gene for the  $\beta$ -lactamase enzyme that can be used as a marker of selection of E. coli cells transformed with plasmid pT7-7(NS5B).

The plasmid pT7-7(NS5B) was then transformed in the E. coli strain BL21(DE53), which is normally employed for high-level expression of genes cloned into expression vectors containing T7 promoter. In this strain of E. coli, the T7 gene polymerase is carried on the bacteriophage  $\lambda$  DE53, which is integrated into the chromosome of BL21 cells (Studier and Moffatt, Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes, (1986), J. Mol. Biol. 189, p. 113-130). Expression from the gene of interest is induced by addition of isopropylthiogalactoside (IPTG) to the growth medium according to a procedure that has been previously described (Studier and Moffatt, 1986). The recombinant NS5B protein fragment containing the RdRp is thus produced in the inclusion bodies of the host cells. Recombinant NS5B protein can be purified from the particulate fraction of E. coli BL21(DE53) extracts and refolded according to procedures that are known in the art (D. R. Thatcher and A. Hitchcock, Protein folding in Biotechnology (1994) in "Mechanism of protein folding" R. H. Pain EDITOR, IRL PRESS, p.229-255). Alternatively, the recombinant NS5B protein could be produced as soluble protein by lowering the temperature of the bacterial growth media below 20°C. The soluble protein could thus be purified from lysates of E. coli substantially as described in Example 5.

#### EXAMPLE 9

Detailed construction of the plasmids in figures

Selected fragments of the cDNA corresponding to the genome of the HCV-BK isolate (HCVBK) were cloned under the strong polyhedrin promoter of the nuclear polyhedrosis virus and flanked by sequences that allowed homologous recombination in a baculovirus vector.

pBac5B contains the HCV-BK sequence comprised between nucleotide 7590 and 9366, and codes for the NS5B protein reported in SEQ ID NO: 1. In order to obtain this plasmid, a cDNA fragment was generated by PCR using synthetic oligonucleotides having the sequences 5'-AAGGATCCATGTCAATGTCCTACACATGGAC-3' (SEQ ID NO: 6) and 5'-AATATTCTGAATTCATCGGTTGGGGAGCAGGTAGATG-3' (SEQ ID NO: 7), respectively. The PCR product was then treated with the Klenow DNA polymerase, digested at the 5'-end with *Bam*HI, and subsequently cloned between the *Bam*HI and *Sma*I sites of the Bluescript SK(+) vector. Subsequently, the cDNA fragment of interest was digested out with the restriction enzymes *Bam*HI and *Hind*III and religated in the same sites of the pBlueBacIII vector (Invitrogen).

pBac25 contains the HCV-BK cDNA region comprised between nucleotides 2759 and 9416 and codes for amino acids 810 to 3010 of the HCV-BK polyprotein (SEQ ID NO: 2). This construct was obtained as follows. First, the 820bp cDNA fragment containing the HCV-BK sequence comprised between nucleotides 2759 and 3578 was obtained from pCD(38-9.4) (Tomei L., Failla, C., Santolini, E., De Francesco, R. and La Monica, N. (1993) NS3 is a Serine Protease Required for Processing of Hepatitis C Virus Polyprotein *J. Virol.*, 67, 4017-4026) by digestion with *Nco*I and cloned in the *Nco*I site of the pBlueBacIII vector (Invitrogen) yielding a plasmid called pBacNCO.. The cDNA fragment containing the HCV-BK sequence comprised between nucleotides 1959 and 9416 was obtained from pCD(38-9.4) (Tomei et al., 1993) by digestion with *Not*I and *Xba*I and cloned in the same sites of the Bluescript SK(+) vector yielding a plasmid called pBlsNX. The cDNA fragment containing the HCV-BK

sequence comprised between nucleotides 3304 and 9416 was obtained from pBlsNX by digestion with *SacII* and *HindIII* and cloned in the same sites of the pBlsNX plasmid, yielding the pBac25 plasmid.

5 pT7-7(DCoH) contains the entire coding region (316 nucleotides) of the rat dimerization cofactor of hepatocyte nuclear factor-1a (DCoH; Mendel, D.B., Khavari, P.A., Conley, P.B., Graves, M.K., Hansen, L.P., Admon, A. and Crabtree, G.R. (1991) Characterization of  
10 a Cofactor that Regulates Dimerization of a Mammalian Homeodomain Protein, *Science* 254, 1762-1767; GenBank accession number: M83740). The cDNA fragment corresponding to the coding sequence for rat DCoH was amplified by PCR using the synthetic oligonucleotide  
15 Dpr1 and Dpr2 that have the sequence TGGCTGGCAAGGCACACAGGCT (SEQ ID NO: 8) and AGGCAGGGTAGATCTATGTC (SEQ ID NO: 9), respectively. The cDNA fragment thus obtained was cloned into the *SmaI* restriction site of the *E. coli* expression vector pT7-7.  
20 The pT7-7 expression vector is a derivative of pBR322 that contains, in addition to the  $\beta$ -lactamase gene and the Col E1 origin of replication, the T7 polymerase promoter  $\phi$ 10 and the translational start site for the T7 gene 10 protein (Tabor S. and Richardson C. C. (1985) A  
25 bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes, *Proc. Natl. Acad. Sci. USA* 82, 1074-1078).

pT7-7(NS5B) contains the HCV sequence from nucleotide 7590 to nucleotide 9366, and codes for the NS5B protein  
30 reported in SEQ ID NO: 1.

In order to obtain this plasmid, a cDNA fragment was generated by PCR using synthetic oligonucleotides having the sequences 5'-TCAATGTCCTACACATGGAC-3' (SEQ ID NO: 10) and 5'-GATCTCTAGATCATCGTTGGGGGAGGAGGTAGATGCC-3' (SEQ ID  
35 NO: 11), respectively. The PCR product was then treated with the Klenow DNA polymerase, and subsequently ligated in the *E. coli* expression vector pT7-7 after linearizing

it with *EcoRI* and blunting its extremities with the Klenow DNA polymerase. Alternatively, cDNA fragment was generated by PCR using synthetic oligonucleotides having the sequences 5'- TGTC AATGTCCTACACATGG-3' (SEQ ID NO: 13) and 5'-AATATT CGAATTCATCGGTTGGGGAGCAGGTAGATG-3' (SEQ ID NO: 14), respectively. The PCR product was then treated with the Klenow DNA polymerase, and subsequently ligated in the *E. coli* expression vector pT7-7 after linearizing it with *NdeI* and blunting its extremities with the Klenow DNA polymerase.



## SEQUENCE LISTING

## GENERAL INFORMATION

- (i) APPLICANT: ISTITUTO DI RICERCHE DI BIOLOGIA  
MOLECOLARE P. ANGELETTI S.p.A.
- 5 (ii) TITLE OF INVENTION: METHOD FOR REPRODUCING  
IN VITRO THE RNA-DEPENDENT RNA POLYMERASE  
AND TERMINAL NUCLEOTIDYL TRANSFERASE  
ACTIVITIES ENCODED BY HEPATITIS C VIRUS  
(HCV)
- 10 (iii) NUMBER OF SEQUENCES: 14
- (iv) CORRESPONDENCE ADDRESS:  
(A)ADDRESSEE: Societa Italiana Brevetti  
(B)STREET: Piazza di Pietra, 39  
(C)CITY: Rome
- 15 (D)COUNTRY: Italy  
(E)POSTAL CODE: 1-00186
- (v) COMPUTER READABLE FORM:  
(A)MEDIUM TYPE: Floppy disk 3.5" 1.44  
MBYTES
- 20 (B)COMPUTER: IBM PC compatible  
(C)OPERATING SYSTEM: PC-DOS/MS-DOS Rev.6.22  
(D)SOFTWARE: Microsoft Word 6.0
- (viii) ATTORNEY INFORMATION  
(A)NAME: DI CERBO, Mario (Dr.)
- 25 (C)REFERENCE: RM/X88530/PCT-DC
- (ix) TELECOMMUNICATION INFORMATION  
(A)TELEPHONE: 06/6785941  
(B)TELEFAX: 06/6794692  
(C)TELEX: 612287 ROPAT
- 30 (1) INFORMATION FOR SEQ ID NO: 1:
- (i) SEQUENCE CHARACTERISTICS  
(A)LENGTH: 591 amino acids  
(B)TYPE: amino acid
- 35 (C)STRANDEDNESS: single  
(D)TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: No  
 (iv) ANTISENSE: No  
 (v) FRAGMENT TYPE: C-terminal fragment  
 (vi) ORIGINAL SOURCE:  
 5 (A) ORGANISM: Hepatitis C Virus  
 (C) ISOLATE : BK  
 (vii) IMMEDIATE SOURCE: cDNA clone pCD(38-9.4)  
 described by Tomei et al. 1993  
 (ix) FEATURE:  
 10 (A) NAME: NS5B Non-structural polyprotein  
 (C) IDENTIFICATION METHOD: Experimentally  
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ser Met Ser Tyr Thr Trp Thr Gly Ala Leu Ile Thr Pro Cys Ala Ala  
 1 5 10 15  
 15 Glu Glu Ser Lys Leu Pro Ile Asn Ala Leu Ser Asn Ser Leu Leu Arg  
 20 25 30  
 His His Asn Met Val Tyr Ala Thr Thr Ser Arg Ser Ala Gly Leu Arg  
 35 40 45  
 Gln Lys Lys Val Thr Phe Asp Arg Leu Gln Val Leu Asp Asp His Tyr  
 20 50 55 60  
 Arg Asp Val Leu Lys Glu Met Lys Ala Lys Ala Ser Thr Val Lys Ala  
 65 70 75 80  
 Lys Leu Leu Ser Val Glu Glu Ala Cys Lys Leu Thr Pro Pro His Ser  
 85 90 95  
 25 Ala Lys Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val Arg Asn Leu Ser  
 100 105 110  
 Ser Lys Ala Val Asn His Ile His Ser Val Trp Lys Asp Leu Leu Glu  
 115 120 125  
 Asp Thr Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn Glu Val  
 30 130 135 140  
 Phe Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala Arg Leu Ile  
 145 150 155 160  
 Val Phe Pro Asp Leu Gly Val Arg Val Cys Glu Lys Met Ala Leu Tyr  
 165 170 175  
 35 Asp Val Val Ser Thr Leu Pro Gln Val Val Met Gly Ser Ser Tyr Gly  
 180 185 190  
 Phe Gln Tyr Ser Pro Gly Gln Arg Val Glu Phe Leu Val Asn Thr Trp

	195	200	205
	Lys Ser Lys Lys Asn Pro Met Gly Phe Ser Tyr Asp Thr Arg Cys Phe		
	210	215	220
5	Asp Ser Thr Val Thr Glu Asn Asp Ile Arg Val Glu Glu Ser Ile Tyr		
	225	230	235 240
	Gln Cys Cys Asp Leu Ala Pro Glu Ala Arg Gln Ala Ile Lys Ser Leu		
	245	250	255
	Thr Glu Arg Leu Tyr Ile Gly Gly Pro Leu Thr Asn Ser Lys Gly Gln		
10	260	265	270
	Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val Leu Thr Thr Ser		
	275	280	285
	Cys Gly Asn Thr Leu Thr Cys Tyr Leu Lys Ala Ser Ala Ala Cys Arg		
	290	295	300
15	Ala Ala Lys Leu Gln Asp Cys Thr Met Leu Val Asn Gly Asp Asp Leu		
	305	310	315 320
	Val Val Ile Cys Glu Ser Ala Gly Thr Gln Glu Asp Ala Ala Ser Leu		
	325	330	335
	Arg Val Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro Pro Gly Asp		
20	340	345	350
	Pro Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser Cys Ser Ser		
	355	360	365
	Asn Val Ser Val Ala His Asp Ala Ser Gly Lys Arg Val Tyr Tyr Leu		
	370	375	380
25	Thr Arg Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala		
	385	390	395 400
	Arg His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile Ile Met Tyr Ala		
	405	410	415
	Pro Thr Leu Trp Ala Arg Met Ile Leu Met Thr His Phe Phe Ser Ile		
30	420	425	430
	Leu Leu Ala Gln Glu Gln Leu Glu Lys Ala Leu Asp Cys Gln Ile Tyr		
	435	440	445
	Gly Ala Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro Gln Ile Ile Glu		
	450	455	460
35	Arg Leu His Gly Leu Ser Ala Phe Ser Leu His Ser Tyr Ser Pro Gly		
	465	470	475 480
	Glu Ile Asn Arg Val Ala Ser Cys Leu Arg Lys Leu Gly Val Pro Pro		

	485	490	495
	Leu Arg Val Trp Arg His Arg Ala Arg Ser Val Arg Ala Arg Leu Leu		
	500	505	510
5	Ser Gln Gly Gly Arg Ala Ala Thr Cys Gly Lys Tyr Leu Phe Asn Trp		
	515	520	525
	Ala Val Lys Thr Lys Leu Lys Leu Thr Pro Ile Pro Ala Ala Ser Arg		
	530	535	540
	Leu Asp Leu Ser Gly Trp Phe Val Ala Gly Tyr Ser Gly Gly Asp Ile		
10	545	550	555
	Tyr His Ser Leu Ser Arg Ala Arg Pro Arg Trp Phe Met Leu Cys Leu		
	565	570	575
	Leu Leu Leu Ser Val Gly Val Gly Ile Tyr Leu Leu Pro Asn Arg		
	580	585	590

(2) INFORMATION FOR SEQ ID NO: 2:

### (i) SEQUENCE CHARACTERISTICS

20 (A) LENGTH: 2201 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear  
(ii) MOLECULE TYPE: polypeptide  
(iii) HYPOTHETICAL: No  
(iv) ANTISENSE: No  
25 (v) FRAGMENT TYPE: C-terminal fragment  
(vii) IMMEDIATE SOURCE: cDNA clone pCD(38-9.4)  
described by Tomei et al. 1993  
(ix) FEATURE:  
(A) NAME: NS2-NS5B Nonstructural Protein  
30 Precursor  
(C) IDENTIFICATION METHOD: Experimentally  
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Asp Arg Glu Met Ala Ala Ser Cys Gly Gly Ala Val Phe Val Gly  
1 5 10 15  
35 Leu Val Leu Leu Thr Leu Ser Pro Tyr Tyr Lys Val Phe Leu Ala Arg  
20 25 30  
Leu Ile Trp Trp Leu Gln Tyr Phe Thr Thr Arg Ala Glu Ala Asp Leu

	35	40	45
	His Val Trp Ile Pro Pro Leu Asn Ala Arg Gly Gly Arg Asp Ala Ile		
	50	55	60
	Ile Leu Leu Met Cys Ala Val His Pro Glu Leu Ile Phe Asp Ile Thr		
5	65	70	75 80
	Lys Leu Leu Ile Ala Ile Leu Gly Pro Leu Met Val Leu Gln Ala Gly		
	85	90	95
	Ile Thr Arg Val Pro Tyr Phe Val Arg Ala Gln Gly Leu Ile His Ala		
	100	105	110
10	Cys Met Leu Val Arg Lys Val Ala Gly Gly His Tyr Val Gln Met Ala		
	115	120	125
	Phe Met Lys Leu Gly Ala Leu Thr Gly Thr Tyr Ile Tyr Asn His Leu		
	130	135	140
	Thr Pro Leu Arg Asp Trp Pro Arg Ala Gly Leu Arg Asp Leu Ala Val		
15	145	150	155 160
	Ala Val Glu Pro Val Val Phe Ser Asp Met Glu Thr Lys Ile Ile Thr		
	165	170	175
	Trp Gly Ala Asp Thr Ala Ala Cys Gly Asp Ile Ile Leu Gly Leu Pro		
	180	185	190
20	Val Ser Ala Arg Arg Gly Lys Glu Ile Leu Leu Gly Pro Ala Asp Ser		
	195	200	205
	Leu Glu Gly Arg Gly Leu Arg Leu Leu Ala Pro Ile Thr Ala Tyr Ser		
	210	215	220
	Gln Gln Thr Arg Gly Leu Leu Gly Cys Ile Ile Thr Ser Leu Thr Gly		
25	225	230	235 240
	Arg Asp Lys Asn Gln Val Glu Gly Glu Val Gln Val Val Ser Thr Ala		
	245	250	255
	Thr Gln Ser Phe Leu Ala Thr Cys Val Asn Gly Val Cys Trp Thr Val		
	260	265	270
30	Tyr His Gly Ala Gly Ser Lys Thr Leu Ala Ala Pro Lys Gly Pro Ile		
	275	280	285
	Thr Gln Met Tyr Thr Asn Val Asp Gln Asp Leu Val Gly Trp Pro Lys		
	290	295	300
	Pro Pro Gly Ala Arg Ser Leu Thr Pro Cys Thr Cys Gly Ser Ser Asp		
35	305	310	315 320
	Leu Tyr Leu Val Thr Arg His Ala Asp Val Ile Pro Val Arg Arg Arg		
	325	330	335



	Val Val Ala Thr Asp Ala Leu Met Thr Gly Tyr Thr Gly Asp Phe Asp	
	625	630 635 640
5	Ser Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr Val Asp Phe Ser	
	645	650 655
	Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Val Pro Gln Asp Ala	
	660	665 670
	Val Ser Arg Ser Gln Arg Arg Gly Arg Thr Gly Arg Gly Arg Arg Gly	
	675	680 685
10	Ile Tyr Arg Phe Val Thr Pro Gly Glu Arg Pro Ser Gly Met Phe Asp	
	690	695 700
	Ser Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr Glu	
	705	710 715 720
	Leu Thr Pro Ala Glu Thr Ser Val Arg Leu Arg Ala Tyr Leu Asn Thr	
15	725	730 735
	Pro Gly Leu Pro Val Cys Gln Asp His Leu Glu Phe Trp Glu Ser Val	
	740	745 750
	Phe Thr Gly Leu Thr His Ile Asp Ala His Phe Leu Ser Gln Thr Lys	
	755	760 765
20	Gln Ala Gly Asp Asn Phe Pro Tyr Leu Val Ala Tyr Gln Ala Thr Val	
	770	775 780
	Cys Ala Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp Gln Met Trp Lys	
	785	790 795 800
	Cys Leu Ile Arg Leu Lys Pro Thr Leu His Gly Pro Thr Pro Leu Leu	
25	805	810 815
	Tyr Arg Leu Gly Ala Val Gln Asn Glu Val Thr Leu Thr His Pro Ile	
	820	825 830
	Thr Lys Tyr Ile Met Ala Cys Met Ser Ala Asp Leu Glu Val Val Thr	
	835	840 845
30	Ser Thr Trp Val Leu Val Gly Gly Val Leu Ala Ala Leu Ala Ala Tyr	
	850	855 860
	Cys Leu Thr Thr Gly Ser Val Val Ile Val Gly Arg Ile Ile Leu Ser	
	865	870 875 880
	Gly Arg Pro Ala Ile Val Pro Asp Arg Glu Leu Leu Tyr Gln Glu Phe	
35	885	890 895
	Asp Glu Met Glu Glu Cys Ala Ser His Leu Pro Tyr Ile Glu Gln Gly	
	900	905 910

	Met	Gln	Leu	Ala	Glu	Gln	Phe	Lys	Gln	Lys	Ala	Leu	Gly	Leu	Leu	Gln
	915							920								925
5	Thr	Ala	Thr	Lys	Gln	Ala	Glu	Ala	Ala	Ala	Pro	Val	Val	Glu	Ser	Lys
	930							935								940
	Trp	Arg	Ala	Leu	Glu	Thr	Phe	Trp	Ala	Lys	His	Met	Trp	Asn	Phe	Ile
	945							950								955
	Ser	Gly	Ile	Gln	Tyr	Leu	Ala	Gly	Leu	Ser	Thr	Leu	Pro	Gly	Asn	Pro
								965								970
10	Ala	Ile	Ala	Ser	Leu	Met	Ala	Phe	Thr	Ala	Ser	Ile	Thr	Ser	Pro	Leu
								980								985
	Thr	Thr	Gln	Ser	Thr	Leu	Leu	Phe	Asn	Ile	Leu	Gly	Gly	Trp	Val	Ala
								995								1000
	Ala	Gln	Leu	Ala	Pro	Pro	Ser	Ala	Ala	Ser	Ala	Phe	Val	Gly	Ala	Gly
15								1010								1015
	Ile	Ala	Gly	Ala	Ala	Val	Gly	Ser	Ile	Gly	Leu	Gly	Lys	Val	Leu	Val
								1025								1030
	Asp	Ile	Leu	Ala	Gly	Tyr	Gly	Ala	Gly	Val	Ala	Gly	Ala	Leu	Val	Ala
								1045								1050
20	Phe	Lys	Val	Met	Ser	Gly	Glu	Met	Pro	Ser	Thr	Glu	Asp	Leu	Val	Asn
								1060								1065
	Leu	Leu	Pro	Ala	Ile	Leu	Ser	Pro	Gly	Ala	Leu	Val	Val	Gly	Val	Val
								1075								1080
	Cys	Ala	Ala	Ile	Leu	Arg	Arg	His	Val	Gly	Pro	Gly	Glu	Gly	Ala	Val
25								1090								1095
	Gln	Trp	Met	Asn	Arg	Leu	Ile	Ala	Phe	Ala	Ser	Arg	Gly	Asn	His	Val
								1105								1110
	Ser	Pro	Thr	His	Tyr	Val	Pro	Glu	Ser	Asp	Ala	Ala	Ala	Arg	Val	Thr
								1125								1130
30	Gln	Ile	Leu	Ser	Ser	Leu	Thr	Ile	Thr	Gln	Leu	Leu	Lys	Arg	Leu	His
								1140								1145
	Gln	Trp	Ile	Asn	Glu	Asp	Cys	Ser	Thr	Pro	Cys	Ser	Gly	Ser	Trp	Leu
								1155								1160
	Arg	Asp	Val	Trp	Asp	Trp	Ile	Cys	Thr	Val	Leu	Thr	Asp	Phe	Lys	Thr
35								1170								1175
	Trp	Leu	Gln	Ser	Lys	Leu	Leu	Pro	Gln	Leu	Pro	Gly	Val	Pro	Phe	Phe
								1185								1190
																1195
																1200



	Ser Cys Gln Arg Gly Tyr Lys Gly Val Trp Arg Gly Asp Gly Ile Met	
	1205	1210 1215
5	Gln Thr Thr Cys Pro Cys Gly Ala Gln Ile Thr Gly His Val Lys Asn	
	1220	1225 1230
	Gly Ser Met Arg Ile Val Gly Pro Lys Thr Cys Ser Asn Thr Trp His	
	1235	1240 1245
	Gly Thr Phe Pro Ile Asn Ala Tyr Thr Thr Gly Pro Cys Thr Pro Ser	
	1250	1255 1260
10	Pro Ala Pro Asn Tyr Ser Arg Ala Leu Trp Arg Val Ala Ala Glu Glu	
	1265	1270 1275 1280
	Tyr Val Glu Val Thr Arg Val Gly Asp Phe His Tyr Val Thr Gly Met	
	1285	1290 1295
	Thr Thr Asp Asn Val Lys Cys Pro Cys Gln Val Pro Ala Pro Glu Phe	
15	1300	1305 1310
	Phe Ser Glu Val Asp Gly Val Arg Leu His Arg Tyr Ala Pro Ala Cys	
	1315	1320 1325
	Arg Pro Leu Leu Arg Glu Glu Val Thr Phe Gln Val Gly Leu Asn Gln	
	1330	1335 1340
20	Tyr Leu Val Gly Ser Gln Leu Pro Cys Glu Pro Glu Pro Asp Val Ala	
	1345	1350 1355 1360
	Val Leu Thr Ser Met Leu Thr Asp Pro Ser His Ile Thr Ala Glu Thr	
	1365	1370 1375
	Ala Lys Arg Arg Leu Ala Arg Gly Ser Pro Pro Ser Leu Ala Ser Ser	
25	1380	1385 1390
	Ser Ala Ser Gln Leu Ser Ala Pro Ser Leu Lys Ala Thr Cys Thr Thr	
	1395	1400 1405
	His His Val Ser Pro Asp Ala Asp Leu Ile Glu Ala Asn Leu Leu Trp	
	1410	1415 1420
30	Arg Gln Glu Met Gly Gly Asn Ile Thr Arg Val Glu Ser Glu Asn Lys	
	1425	1430 1435 1440
	Val Val Val Leu Asp Ser Phe Asp Pro Leu Arg Ala Glu Glu Asp Glu	
	1445	1450 1455
	Arg Glu Val Ser Val Pro Ala Glu Ile Leu Arg Lys Ser Lys Lys Phe	
35	1460	1465 1470
	Pro Ala Ala Met Pro Ile Trp Ala Arg Pro Asp Tyr Asn Pro Pro Leu	
	1475	1480 1485

-32-

	Leu Glu Ser Trp Lys Asp Pro Asp Tyr Val Pro Pro Val Val His Gly
	1490 1495 1500
5	Cys Pro Leu Pro Pro Ile Lys Ala Pro Pro Ile Pro Pro Pro Arg Arg
	1505 1510 1515 1520
	Lys Arg Thr Val Val Leu Thr Glu Ser Ser Val Ser Ser Ala Leu Ala
	1525 1530 1535
	Glu Leu Ala Thr Lys Thr Phe Gly Ser Ser Glu Ser Ser Ala Val Asp
	1540 1545 1550
10	Ser Gly Thr Ala Thr Ala Leu Pro Asp Gln Ala Ser Asp Asp Gly Asp
	1555 1560 1565
	Lys Gly Ser Asp Val Glu Ser Tyr Ser Ser Met Pro Pro Leu Glu Gly
	1570 1575 1580
	Glu Pro Gly Asp Pro Asp Leu Ser Asp Gly Ser Trp Ser Thr Val Ser
15	1585 1590 1595 1600
	Glu Glu Ala Ser Glu Asp Val Val Cys Cys Ser Met Ser Tyr Thr Trp
	1605 1610 1615
	Thr Gly Ala Leu Ile Thr Pro Cys Ala Ala Glu Glu Ser Lys Leu Pro
	1620 1625 1630
20	Ile Asn Ala Leu Ser Asn Ser Leu Leu Arg His His Asn Met Val Tyr
	1635 1640 1645
	Ala Thr Thr Ser Arg Ser Ala Gly Leu Arg Gln Lys Lys Val Thr Phe
	1650 1655 1660
	Asp Arg Leu Gln Val Leu Asp Asp His Tyr Arg Asp Val Leu Lys Glu
25	1665 1670 1675 1680
	Met Lys Ala Lys Ala Ser Thr Val Lys Ala Lys Leu Leu Ser Val Glu
	1685 1690 1695
	Glu Ala Cys Lys Leu Thr Pro Pro His Ser Ala Lys Ser Lys Phe Gly
	1700 1705 1710
30	Tyr Gly Ala Lys Asp Val Arg Asn Leu Ser Ser Lys Ala Val Asn His
	1715 1720 1725
	Ile His Ser Val Trp Lys Asp Leu Leu Glu Asp Thr Val Thr Pro Ile
	1730 1735 1740
	Asp Thr Thr Ile Met Ala Lys Asn Glu Val Phe Cys Val Gln Pro Glu
35	1745 1750 1755 1760
	Lys Gly Gly Arg Lys Pro Ala Arg Leu Ile Val Phe Pro Asp Leu Gly
	1765 1770 1775

SUBSTITUTE SHEET (RULE 26)

	Val Arg Val Cys Glu Lys Met Ala Leu Tyr Asp Val Val Ser Thr Leu	
	1780	1785 1790
5	Pro Gln Val Val Met Gly Ser Ser Tyr Gly Phe Gln Tyr Ser Pro Gly	
	1795	1800 1805
	Gln Arg Val Glu Phe Leu Val Asn Thr Trp Lys Ser Lys Lys Asn Pro	
	1810	1815 1820
	Met Gly Phe Ser Tyr Asp Thr Arg Cys Phe Asp Ser Thr Val Thr Glu	
	1825	1830 1835 1840
10	Asn Asp Ile Arg Val Glu Glu Ser Ile Tyr Gln Cys Cys Asp Leu Ala	
	1845	1850 1855
	Pro Glu Ala Arg Gln Ala Ile Lys Ser Leu Thr Glu Arg Leu Tyr Ile	
	1860	1865 1870
15	Gly Gly Pro Leu Thr Asn Ser Lys Gly Gln Asn Cys Gly Tyr Arg Arg	
	1875	1880 1885
	Cys Arg Ala Ser Gly Val Leu Thr Thr Ser Cys Gly Asn Thr Leu Thr	
	1890	1895 1900
	Cys Tyr Leu Lys Ala Ser Ala Ala Cys Arg Ala Ala Lys Leu Gln Asp	
	1905	1910 1915 1920
20	Cys Thr Met Leu Val Asn Gly Asp Asp Leu Val Val Ile Cys Glu Ser	
	1925	1930 1935
	Ala Gly Thr Gln Glu Asp Ala Ala Ser Leu Arg Val Phe Thr Glu Ala	
	1940	1945 1950
	Met Thr Arg Tyr Ser Ala Pro Pro Gly Asp Pro Pro Gln Pro Glu Tyr	
25	1955	1960 1965
	Asp Leu Glu Leu Ile Thr Ser Cys Ser Ser Asn Val Ser Val Ala His	
	1970	1975 1980
	Asp Ala Ser Gly Lys Arg Val Tyr Tyr Leu Thr Arg Asp Pro Thr Thr	
	1985	1990 1995 2000
30	Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg His Thr Pro Val Asn	
	2005	2010 2015
	Ser Trp Leu Gly Asn Ile Ile Met Tyr Ala Pro Thr Leu Trp Ala Arg	
	2020	2025 2030
	Met Ile Leu Met Thr His Phe Phe Ser Ile Leu Leu Ala Gln Glu Gln	
35	2035	2040 2045
	Leu Glu Lys Ala Leu Asp Cys Gln Ile Tyr Gly Ala Cys Tyr Ser Ile	
	2050	2055 2060

Glu Pro Leu Asp Leu Pro Gln Ile Ile Glu Arg Leu His Gly Leu Ser  
 2065                      2070                      2075                      2080

Ala Phe Ser Leu His Ser Tyr Ser Pro Gly Glu Ile Asn Arg Val Ala  
 5                      2085                      2090                      2095

Ser Cys Leu Arg Lys Leu Gly Val Pro Pro Leu Arg Val Trp Arg His  
                     2100                      2105                      2110

Arg Ala Arg Ser Val Arg Ala Arg Leu Leu Ser Gln Gly Gly Arg Ala  
                     2115                      2120                      2125

10    Ala Thr Cys Gly Lys Tyr Leu Phe Asn Trp Ala Val Lys Thr Lys Leu  
                     2130                      2135                      2140

Lys Leu Thr Pro Ile Pro Ala Ala Ser Arg Leu Asp Leu Ser Gly Trp  
 2145                      2150                      2155                      2160

Phe Val Ala Gly Tyr Ser Gly Gly Asp Ile Tyr His Ser Leu Ser Arg  
 15                      2165                      2170                      2175

Ala Arg Pro Arg Trp Phe Met Leu Cys Leu Leu Leu Leu Ser Val Gly  
                     2180                      2185                      2190

Val Gly Ile Tyr Leu Leu Pro Asn Arg  
                     2195                      2200

20    (3)    INFORMATION FOR SEQ ID NO: 3

          (i)    SEQUENCE CHARACTERISTICS

                    (A) LENGTH: 26 nucleotides

                    (B) TYPE: nucleic acid

25                      (C) STRANDEDNESS: single

                    (D) TOPOLOGY: linear

          (ii)    MOLECULE TYPE: synthetic DNA

          (iii)    HYPOTHETICAL: No

          (iv)    ANTISENSE: No

30                      (vii) IMMEDIATE SOURCE: oligonucleotide

                                    synthesizer

          (ix)    FEATURE:

                    (A) NAME: oligo a

                    (C) IDENTIFICATION METHOD: Polyacrylamide

35                                      gel

          (xi)    SEQUENCE DESCRIPTION: SEQ ID NO: 3

GCCGAGATGC CATCTTCAA CAGTTC

26

## (4) INFORMATION FOR SEQ ID NO: 4

- 5 (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 24 nucleotides  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: synthetic DNA  
(iii) HYPOTHETICAL: No  
(iv) ANTISENSE: No  
(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer
- 15 (ix) FEATURE:  
(A) NAME: oligo b  
(C) IDENTIFICATION METHOD: Polyacrylamide  
gel
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4

20 GTGTACAACA AGGTCCATAT CACC

24

## (5) INFORMATION FOR SEQ ID NO: 5

- 25 (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 24 nucleotides  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- 30 (ii) MOLECULE TYPE: synthetic DNA  
(iii) HYPOTHETICAL: No  
(iv) ANTISENSE: No  
(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer
- 35 (ix) FEATURE:  
(A) NAME: oligo c  
(C) IDENTIFICATION METHOD: Polyacrylamide  
gel
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5

GGTCTTTCTG AACGGGATAT AAAC

24

- (6) INFORMATION FOR SEQ ID NO: 6:
- 5 (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 31 nucleotides  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- 10 (ii) MOLECULE TYPE: synthetic DNA  
(iii) HYPOTHETICAL: No  
(iv) ANTISENSE: No  
(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer
- 15 (ix) FEATURE:  
(A) NAME: 5'-5B  
(C) IDENTIFICATION METHOD: Polyacrylamide  
gel
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6

20

AAGGATCCAT GTCAATGTCC TACACATGGA C

31

- (7) INFORMATION FOR SEQ ID NO: 7:
- 25 (i) SEQUENCE CHARACTERISTICS  
(A) LENGTH: 36 nucleotides  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: synthetic DNA
- 30 (iii) HYPOTHETICAL: No  
(iv) ANTISENSE: Yes  
(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer
- (ix) FEATURE:  
35 (A) NAME: 3'-5B  
(C) IDENTIFICATION METHOD: Polyacrylamide  
gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7

AATATTCGAA TTCATCGGTT GGGGAGCAGG TAGATG

36

5 (8) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 22 nucleotides

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTISENSE: No

15 (vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer

(ix) FEATURE:

(A) NAME: Dpr1

(C) IDENTIFICATION METHOD: Polyacrylamide  
gel

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8

TGGCTGGCAA GGCACACAGG CT

22

(9) INFORMATION FOR SEQ ID NO: 9

25 (i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 20 nucleotides

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTISENSE: Yes

(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer

35 (ix) FEATURE:

(A) NAME: Dpr2

(C) IDENTIFICATION METHOD: Polyacrylamide  
gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9

5 AGGCAGGGTA GATCTATGTC

20

(10) INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 20 nucleotides

10 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: No

15 (iv) ANTISENSE: No

(vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer

(ix) FEATURE:

(A) NAME: NS5B-5' (1)

20 (C) IDENTIFICATION METHOD: Polyacrylamide  
gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10

TCAATGTCCT ACACATGGAC

20

25

(11) INFORMATION FOR SEQ ID NO: 11

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 38 nucleotides

(B) TYPE: nucleic acid

30 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTISENSE: Yes

35 (vii) IMMEDIATE SOURCE: oligonucleotide  
synthesizer

(ix) FEATURE:



(A)NAME: HCVA-13

(C)IDENTIFICATION METHOD: Polyacrylamide  
gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11

5

GATCTCTAGA TCATCGGTTG GGGGAGGAGG TAGATGCC

38

(12) INFORMATION FOR SEQ ID NO: 12

(i) SEQUENCE CHARACTERISTICS

10

(A)LENGTH: 399 nucleotides

(B)TYPE: nucleic acid

(C)STRANDEDNESS: single

(D)TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

15

(iii) HYPOTHETICAL: No

(iv) ANTISENSE: No

(vi) ORIGINAL SOURCE:

(A)ORGANISM: Rattus Norvegicus

(B)STRAIN : Sprague-Dawley

20

(vii) IMMEDIATE SOURCE: pT7-7(DCoH)

(ix) FEATURE:

(A)NAME: D-RNA

(C)IDENTIFICATION METHOD: Polyacrylamide  
gel

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12

GGGAGACCAC AACGGUUUCC CUCUAGAAU AAUUUUGUUU AACUUUAAGA AGGAGAUUA 60

CAUAUGGCUA GAAUUCGCGC CCUGGCUGGC AAGGCACACA GGCUGAGUGC UGAGGAACGG 120

GACCAGCUGC UGCCAAACCU GCGGGCCGUG GGGUGGAAUG AACUGGAAGG CCGAGAUGCC 180

30 AUCUCAAAC AGUCCAUUU UAAAGACUUC AACAGGGCUU UUGGCUUCAU GACAAGAGUC 240

GCCCUGCAGG CUGAAAAGCU GGACCACCAU CCCGAGUGGU UUAACGUGUA CAACAAGGUC 300

CAUAUCACCU UGAGCACCCA CGAUGUGCC GGUCUUUCUG AACGGGAUUA AAACCUGGCC 360

AGCUUCAUCG AACAAGUUGC CGUGUCUAUG ACAUAGAUC 399

(13) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 20 nucleotides

(B) TYPE: nucleic acid

5 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTISENSE: No

10 (vii) IMMEDIATE SOURCE: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME: NS5B-up

(C) IDENTIFICATION METHOD: Polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13

15

TGTCAATGTC CTACACATGG

20

(14) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS

20 (A) LENGTH: 38 nucleotides

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

25 (iii) HYPOTHETICAL: No

(iv) ANTISENSE: Yes

(vii) IMMEDIATE SOURCE: oligonucleotide synthesizer

(ix) FEATURE:

(A) NAME: 3'-5B

30 (C) IDENTIFICATION METHOD: Polyacrylamide gel

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14

AATATTCGAA TTCATCGGTT GGGGAGCAGG TAGATG

36

CLAIMS

1. A method for reproducing in vitro the RNA-dependent RNA polymerase activity or the terminal nucleotidyl transferase activity encoded by hepatitis C virus, characterized in that sequences containing NS5B (SEQ ID NO: 1) are used in the reaction mixture.

2. The method for reproducing in vitro the RNA-dependent RNA polymerase activity encoded by HCV according to claim 1, in which NS5B is incorporated in the reaction mixture as NS2-NS5B precursor, said precursor generating, by means of multiple proteolytic events that occur in the overproducing organism, an enzymatically active form of NS5B.

3. The method for reproducing in vitro the terminal nucleotidyl transferase activity encoded by HCV according to claim 1, in which NS5B is incorporated in the reaction mixture as NS2-NS5B precursor, said precursor generating, by means of multiple proteolytic events that occur in the overproducing organism, an enzymatically active form of NS5B.


4. A composition of matter, characterized in that it contains NS5B sequences according to claims 1 to 3.

5. A composition of matter according to claim 4, comprising the proteins whose sequences are described in SEQ ID NO: 1, in sequences contained therein or derived therefrom.

6. Use of the compositions of matter according to claims 4 and 5 to set up an enzymatic test capable of selecting, for therapeutic purposes, compounds that inhibit the enzymatic activity associated with NS5B.

7. Method for reproducing in vitro the RNA-dependent RNA polymerase and terminal nucleotidyl transferase activities of NS5B, compositions of matter and use of said compositions of matter to set up an enzymatic test capable of selecting, for therapeutic purposes, compounds that inhibit the enzymatic activities associated with NS5B, according to the above description, examples and claims.

1/3



P ETL = promoter of the gene coding for the PCNA protein

P PH = promoter of the polyhedrin gene


Amp = gene coding for the β-lactamase enzyme  
(ampicillin resistance)

LacZ (β-gal) = gene coding for the β-galactosidase enzyme


Col E1 = pBR322 replication origin

FIG. 1

2/3  
pT7-7(DCoH)



pT7-7(NS5B)



Ø10 = bacteriophage T7 Ø10 promoter

rbs = Shine-Dalgarno ribosome binding site

ATG = translation initiation site of the protein  
coded by the bacteriophage T7 gene 10

β-lactamase = gene coding for the β-lactamase enzyme  
(ampicillin resistance)

Col E1 = pBR322 replication origin

FIG.2




FIG.3

## INTERNATIONAL SEARCH REPORT

Patent Application No.

PCT/IT 96/00106

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/54 C12N9/12 C12Q1/48 G01N33/573

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 463 848 (UNIV OSAKA RES FOUND) 2 January 1992	4,5
Y	see page 3, line 45 - line 50 see page 11, line 7 see page 19, line 39 - line 46 see page 21, line 1 - page 29, line 42 see page 50, line 26 - page 53, line 25 see claims 1-21; figure 1 ---	1,2,6,7
X	EP,A,0 464 287 (UNIV OSAKA RES FOUND) 8 January 1992	4,5
Y	see page 11, line 13 - page 16, line 45; claims 1-31 --- -/--	1,2,6,7

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

29 August 1996

Date of mailing of the international search report

03.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

## INTERNATIONAL SEARCH REPORT

Inter national Patent No

PCT/IT 99/00106

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. VIROLOGY, vol. 67, no. 7, July 1993, AM.SOC.MICROBIOL.,WASHINGTON,US, pages 4017-4026, XP000601449 L. TOMEI ET AL.: "NS3 is a serine protease required for processing of hepatitis C virus polyprotein" cited in the application	4,5
Y	see page 4017, right-hand column, line 18 - line 22 see page 4018, right-hand column, line 35 - page 4019, left-hand column, line 2 see page 4021, left-hand column, line 30 - line 35	1,2,6,7
Y	--- WO,A,94 05809 (FOX CHASE CANCER CENTER ;SEEGER CHRISTOPH (US)) 17 March 1994 see the whole document ---	1,2,6,7
A	--- J. VIROLOGY, vol. 68, no. 8, August 1994, AM.SOC.MICROBIOL.,WASHINGTON,US, pages 5045-5055, XP002012001 R. BARTENSCHLAGER ET AL.: "Kinetic and structural analysis of hepatitis C virus polyprotein processing" see the whole document ---	1-7
A	--- J. VIROLOGY, vol. 68, no. 12, December 1994, AM.SOC.MICROBIOL.,WASHINGTON,US, pages 8147-8157, XP002012002 C LIN ET AL.: "Hepatitis C virus NS3 serine proteinase: trans-cleavage requirements and processing kinetics" see the whole document ---	1-7
A	--- PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 87, no. 6, 1 March 1990, pages 2057-2061, XP000310581 MILLER R H ET AL: "HEPATITIS C VIRUS SHARES AMINO ACID SEQUENCE SIMILARITY WITH PESTIVIRUSES AND FLAVIVIRUSES AS WELL AS MEMBERS OF TWO PLANT VIRUS SUPERGROUPS" see the whole document ---	1-7
P,X	--- WO,A,95 22985 (ISTITUTO DI RICERCHE DI BIOLOG ;FRANCESCO RAFFAELE DE (IT); FAILLA) 31 August 1995	4,5
P,Y	see the whole document ---	1,2,6,7
	---	

-/--



## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IT 96/00106

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>EMBO (EUROPEAN MOLECULAR BIOLOGY ORGANIZATION) JOURNAL 15 (1). 1996. 12-22. ISSN: 0261-4189, January 1996, XP002012003 BEHRENS S-E ET AL: "Identification and properties of the RNA-dependent RNA polymerase of hepatitis C virus." see the whole document -----</p>	1-7

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IT 97/00106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0463848	02-01-92	AU-B- 652941	15-09-94
		AU-B- 7925691	02-01-92
		CA-A- 2045326	26-12-91
		JP-A- 7258290	09-10-95
		CN-A- 1059758	25-03-92
		AU-B- 650634	30-06-94
		AU-B- 6860891	02-01-92
		CA-A- 2045323	26-12-91
		CN-A- 1057861	15-01-92
		EP-A- 0464287	08-01-92
		JP-A- 6303980	01-11-94
		KR-B- 9512971	24-10-95
EP-A-0464287	08-01-92	AU-B- 650634	30-06-94
		AU-B- 6860891	02-01-92
		AU-B- 652941	15-09-94
		AU-B- 7925691	02-01-92
		CA-A- 2045323	26-12-91
		CA-A- 2045326	26-12-91
		CN-A- 1057861	15-01-92
		CN-A- 1059758	25-03-92
		EP-A- 0463848	02-01-92
		JP-A- 6303980	01-11-94
		JP-A- 7258290	09-10-95
		KR-B- 9512971	24-10-95
WO-A-9405809	17-03-94	US-A- 5360714	01-11-94
		US-A- 5334525	02-08-94
		AU-B- 5095793	29-03-94
		CA-A- 2143194	17-03-94
		EP-A- 0673437	27-09-95
		JP-T- 8503846	30-04-96
WO-A-9522985	31-08-95	AU-B- 1822395	11-09-95

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**